ELSEVIER

Molecular Brain Research 24 (1994) 357–358

MOLECULAR BRAIN RESEARCH

Author Index

Abdou, A., see Greenwood, R.S. (24) 20
Abe-Dohmae, S., Tanaka, R. and Harada, N.
Cell type- and region-specific expression of aromatase mRNA in cultured brain cells (24) 153
Akagi, H., see Uchiyama, M. (24) 295
Almazan, G., see Radhakrishna, M. (24) 118
Arnold, A.P., see Shen, P. (24) 227
Ashcroft, F.M., see Kenna, S. (24) 353
Ashcroft, S.J.H., see Kenna, S. (24) 353

Bakalkin, G., Yakovleva, T. and Terenius, L.
Prodynorphin gene expression relates to NF-κB factors (24) 301
Bale, T.L., see Szot, P. (24) 1
Banerjee, S.A., Roffler-Tarlov, S., Szabo, M., Frohman, L. and Chikaraishi, D.M.
DNA regulatory sequences of the rat tyrosine hydroxylase gene direct correct catecholaminergic cell-type specificity of a human growth hormone reporter in the CNS of transgenic mice causing a dwarf phenotype (24) 89
Barnes Jr., E.M., see Jalilian Tehrani, M.H. (24) 55
Baudet, C., see Neveu, I. (24) 70
Baudier, J., see Palacios, G. (24) 107
Bonell, M.A., see Wu, P. (24) 27
Boone Jr., J.B. and McMillen, D.
Proenkephalin gene expression is altered in the brain of spontaneously hypertensive rats during the development of hypertension (24) 320
Brachet, P., see Neveu, I. (24) 70
Brady, D.R., see Chandrasekaran, K. (24) 336
Brown, I.R., see Mendis, D.B. (24) 11

Cagampang, F.R.A., Yang, J., Nakayama, Y., Fukuhara, C. and Inouye, S.-I.T.
Circadian variation of arginine-vasopressin messenger RNA in the rat suprachiasmatic nucleus (24) 179
Campagnoni, A.T., see Shen, P. (24) 227
Campagnoni, C.W., see Shen, P. (24) 227
Cashman, N.R., see Ledoux, S. (24) 140
Chandrasekaran, K., Giordano, T., Brady, D.R., Stoll, J., Martin, L.J. and Rapoport, S.I.
Impairment in mitochondrial cytochrome oxidase gene expression in Alzheimer disease (24) 336
Chikaraishi, D.M., see Banerjee, S.A. (24) 89
Cho, J.Y., see Huh, S.O. (24) 145
Christakos, S., see Lee, S. (24) 34

De Blas, A.L., see Miralles, C.P. (24) 129
De Luca, H.F., see Neveu, I. (24) 70
Devinsky, O., see Lee, S. (24) 34
Dorsa, D.M., see Szot, P. (24) 1
Dragunow, M., see Hughes, P. (24) 166
Dressler, H., see Kelley, K.A. (24) 261
Drinkwater, D., see Kelley, K.A. (24) 261
Dubner, R., see Gu, J. (24) 77

Federoff, H.J., see Geschwind, M.D. (24) 327
Fletcher, D.L., see Zoeller, R.T. (24) 185
Friedrich Jr., V.L., see Kelley, K.A. (24) 261
Frohman, L., see Banerjee, S.A. (24) 89
Fukuda, K., see Houtani, T. (24) 347
Fukuhara, C., see Cagampang, F.R.A. (24) 179

Geller, A.I., see Geschwind, M.D. (24) 327
Geschwind, M.D., Kessler, J.A., Geller, A.I. and Federoff, H.J.
Transfer of the nerve growth factor gene into cell lines and cultured neurons using a defective Herpes Simplex virus vector. Transfer of the NGF gene into cells by a HSV-1 vector (24) 327
Giordano, T., see Chandrasekaran, K. (24) 336
Greenwood, R.S., Meeker, R.B., Abdou, A. and Hayward, J.N.
Kindled seizures induce a long-term increase in vasopressin mRNA (24) 20
Grouzmann, E., see Wu, P. (24) 27
Gu, J., Ren, K., Dubner, R. and Iadarola, M.J.
Cloning of a DNA binding protein that is a tyrosine kinase substrate and recognizes an upstream initiator-like sequence in the promoter of the preprodynorphin gene (24) 77
Gutiérrez, A., see Miralles, C.P. (24) 129

Harada, N., see Abe-Dohmae, S. (24) 153
Hayward, J.N., see Greenwood, R.S. (24) 20
Hebert, S., see Kenna, S. (24) 353
Herrup, K., see Kelley, K.A. (24) 261
Hirai, K., see Uchiyama, M. (24) 295
Hishinuma, F., see Uchiyama, M. (24) 295
Ho, K., see Kenna, S. (24) 353
Houtani, T., Ueyama, T., Takeshima, H., Kato, S., Fukuda, K., Mori, K. and Sugimoto, T.
μ Opioid receptor: expression and vagotomy-induced depletion of the mRNA in medullary preganglionic neurons (24) 347
Howells, R., see Lee, S. (24) 34
Hu, Y., see Kelley, K.A. (24) 261
Hughes, P. and Dragunow, M.
Activation of pirenzepine-sensitive muscarinic receptors induces a specific pattern of immediate-early gene expression in rat brain neurons (24) 166
Huh, S.O., Park, D.H., Cho, J.Y., Joh, T.H. and Son, J.H.
A 6.1 kb 5′ upstream region of the mouse tryptophan hydroxylase gene directs expression of *E. coli lacZ* to major serotonergic brain regions and pineal gland in transgenic mice (24) 145

Iadarola, M.J., see Gu, J. (24) 77
Imai, Y., Suzuki, Y., Tohyama, M., Wanaka, A. and Takagi, T.
Cloning and expression of a neural differentiation-associated gene, p205, in the embryonal carcinoma cell line P19 and in the developing mouse (24) 313
Inouye, S.-I.T., see Cagampang, F.R.A. (24) 179

Jalilian Tehrani, M.H. and Barnes Jr., E.M.
$GABA_A$ receptors in mouse cortical homogenates are phosphorylated by endogenous protein kinase A (24) 55
Jehan, F., see Neveu, I. (24) 70
Joh, T.H., see Huh, S.O. (24) 145
Jope, R.S., see Unlap, T. (24) 275
Junier, M.P., Suzuki, F., Onteniente, B. and Peschanski, M.
Target-deprived CNS neurons express the NGF gene while reactive glia around their axonal terminals contain low and high affinity NGF receptors (24) 247

Kampf, K., see Shen, P. (24) 227
Kato, S., see Houtani, T. (24) 347
Kawai, Y., see Senba, E. (24) 283
Kelley, K.A., Friedrich Jr., V.L., Sonshine, A., Hu, Y., Lax, J., Li, J., Drinkwater, D., Dressler, H. and Herrup, K.
Expression of Thy-1/*lacZ* fusion genes in the CNS of transgenic mice (24) 261
Kenna, S., Röper, J., Ho, K., Hebert, S., Ashcroft, S.J.H. and Ashcroft, F.M.
Differential expression of the inwardly-rectifying K-channel ROMK1 in rat brain (24) 353
Kessler, J.A., see Geschwind, M.D. (24) 327
Khan, Z.U., see Miralles, C.P. (24) 129

Lasoń, W. and Przewłocki, R.
Seizure-induced expression of G proteins in the rat hippocampus (24) 65

Lax, J., see Kelley, K.A. (24) 261
Ledoux, S., Nalbantoglu, J. and Cashman, N.R.
Amyloid precursor protein gene expression in neural cell lines: influence of DNA cytosine methylation (24) 140
Lee, S., Miskovsky, J., Williamson, J., Howells, R., Devinsky, O., Lothman, E. and Christakos, S.
Changes in glutamate receptor and proenkephalin gene expression after kindled seizures (24) 34
Lee, S., Rivier, C. and Torres, G.
Induction of c-*fos* and CRF mRNA by MK-801 in the parvocellular paraventricular nucleus of the rat hypothalamus (24) 192
Li, C., see Schinkmann, K. (24) 238
Li, J., see Kelley, K.A. (24) 261
Lothman, E., see Lee, S. (24) 34
Luque, J.M., Malherbe, P. and Richards, J.G.
Localization of $GABA_A$ receptor subunit mRNAs in the rat locus coeruleus (24) 219

Malherbe, P., see Luque, J.M. (24) 219
Martin, L.J., see Chandrasekaran, K. (24) 336
McMillen, D., see Boone Jr., J.B. (24) 320
Meeker, R.B., see Greenwood, R.S. (24) 20
Mendis, D.B. and Brown, I.R.
Expression of the gene encoding the extracellular matrix glycoprotein SPARC in the developing and adult mouse brain (24) 11
Mengod, G., see Palacios, G. (24) 107
Meyer, E.M., see Wu, P. (24) 27
Mhatre, M.C. and Ticku, M.K.
Chronic GABA treatment downregulates the $GABA_A$ receptor α_2 and α_3 subunit mRNAS as well as polypeptide expression in primary cultured cerebral cortical neurons (24) 159
Millard, W.J., see Wu, P. (24) 27
Miralles, C.P., Gutiérrez, A., Khan, Z.U., Vitorica, J. and De Blas, A.L.
Differential expression of the short and long forms of the γ_2 subunit of the $GABA_A$/benzodiazepine receptors (24) 129
Miskovsky, J., see Lee, S. (24) 34
Mori, K., see Houtani, T. (24) 347

Nakayama, Y., see Cagampang, F.R.A. (24) 179
Nalbantoglu, J., see Ledoux, S. (24) 140
Naveilhan, P., see Neveu, I. (24) 70
Neveu, I., Naveilhan, P., Jehan, F., Baudet, C., Wion, D., De Luca, H.F. and Brachet, P.
1,25-Dihydroxyvitamin D_3 regulates the synthesis of nerve growth factor in primary cultures of glial cells (24) 70
Nishizaki, T. and Sumikawa, K.
A cAMP-dependent Ca^{2+} signalling pathway at the endplate provided by the γ to ϵ subunit switch in ACh receptors (24) 341
Noguchi, K., see Senba, E. (24) 283

Onteniente, B., see Junier, M.P. (24) 247

Palacios, G., Mengod, G., Sarasa, M., Baudier, J. and Palacios, J.M.
De novo synthesis of GAP-43: in situ hybridization histochemistry and light and electron microscopy immunocytochemical studies in regenerating motor neurons of cranial nerve nuclei in the rat brain (24) 107
Palacios, J.M., see Palacios, G. (24) 107
Palacios, J.M., see Waeber, C. (24) 199
Park, D.H., see Huh, S.O. (24) 145
Peschanski, M., see Junier, M.P. (24) 247
Przewłocki, R., see Lasoń, W. (24) 65

Radhakrishna, M. and Almazan, G.
Protein kinases mediate basic fibroblast growth factor's stimulation of proliferation and c-*fos* induction in oligodendrocyte progenitors (24) 118
Rapoport, S.I., see Chandrasekaran, K. (24) 336
Ren, K., see Gu, J. (24) 77
Richards, J.G., see Luque, J.M. (24) 219
Rivier, C., see Lee, S. (24) 192
Robinson, G.A.
Immediate early gene expression in axotomized and regenerating retinal ganglion cells of the adult rat (24) 43
Roffler-Tarlov, S., see Banerjee, S.A. (24) 89
Röper, J., see Kenna, S. (24) 353

Sarasa, M., see Palacios, G. (24) 107
Schinkmann, K. and Li, C.
Comparison of two *Caenorhabditis* genes encoding FMRFamide(Phe-Met-Arg-Phe-NH_2)-like peptides (24) 238
Schlinger, B.A., see Shen, P. (24) 227
Schwartz, J.H., see Sossin, W.S. (24) 210
Senba, E., Umemoto, S., Kawai, Y. and Noguchi, K.
Differential expression of *fos* family and *jun* family mRNAs in the rat hypothalamo-pituitary-adrenal axis after immobilization stress (24) 283
Shen, P., Campagnoni, C.W., Kampf, K., Schlinger, B.A., Arnold, A.P. and Campagnoni, A.T.
Isolation and characterization of a zebra finch aromatase cDNA: in situ hybridization reveals high aromatase expression in brain (24) 227
Son, J.H., see Huh, S.O. (24) 145
Sonshine, A., see Kelley, K.A. (24) 261
Sossin, W.S. and Schwartz, J.H.
Translocation of protein kinase Cs in *Aplysia* neurons: evidence for complex regulation (24) 210
Stoll, J., see Chandrasekaran, K. (24) 336
Sugimoto, T., see Houtani, T. (24) 347
Sumikawa, K., see Nishizaki, T. (24) 341
Suzuki, F., see Junier, M.P. (24) 247
Suzuki, Y., see Imai, Y. (24) 313
Szabo, M., see Banerjee, S.A. (24) 89
Szot, P., Bale, T.L. and Dorsa, D.M.
Distribution of messenger RNA for the vasopressin V1a receptor in the CNS of male and female rats (24) 1

Takagi, T., see Imai, Y. (24) 313
Takeshima, H., see Houtani, T. (24) 347
Tanaka, R., see Abe-Dohmae, S. (24) 153
Terenius, L., see Bakalkin, G. (24) 301
Ticku, M.K., see Mhatre, M.C. (24) 159
Tohyama, M., see Imai, Y. (24) 313
Torres, G., see Lee, S. (24) 192

Uchiyama, M., Hirai, K., Hishinuma, F. and Akagi, H.
Down-regulation of glycine receptor channels by protein kinase C in *Xenopus* oocytes injected with synthetic RNA (24) 295
Ueyama, T., see Houtani, T. (24) 347
Umemoto, S., see Senba, E. (24) 283
Unlap, T. and Jope, R.S.
Dexamethasone attenuates kainate-induced AP-1 activation in rat brain (24) 275

Vitorica, J., see Miralles, C.P. (24) 129

Waeber, C. and Palacios, J.M.
Binding sites for 5-hydroxytryptamine-2 receptor agonists are predominantly located in striosomes in the human basal ganglia (24) 199
Wanaka, A., see Imai, Y. (24) 313
Williamson, J., see Lee, S. (24) 34
Wion, D., see Neveu, I. (24) 70
Wu, P., Ziska, D., Bonell, M.A., Grouzmann, E., Millard, W.J. and Meyer, E.M.
Differential neuropeptide Y gene expression in post-mitotic versus dividing neuroblastoma cells driven by an adeno-associated virus vector (24) 27

Yakovleva, T., see Bakalkin, G. (24) 301
Yang, J., see Cagampang, F.R.A. (24) 179

Ziska, D., see Wu, P. (24) 27
Zoeller, R.T. and Fletcher, D.L.
A single administration of ethanol simultaneously increases c-*fos* mRNA and reduces c-*jun* mRNA in the hypothalamus and hippocampus (24) 185